

A class of multidimensional quadratic BSDEs

Zhongmin Qian^{*}, Yimin Yang[†] and Shujin Wu[‡]

March 14, 2017

Abstract

In this paper we study a multidimensional quadratic BSDE with a particular class of product generators and bounded terminal values, and give a result of existence of solution in a suitable Banach space under some constraints on parameters.

Key words: Multidimensional BSDEs, quadratic BSDEs, BMO martingales.

1 Introduction

A multidimensional quadratic Backward Stochastic Differential Equation (BSDE) on $[0, T]$ with T being the terminal time, according to the formulation put forward by Pardoux and Peng [14], is a stochastic integral equation with

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s \quad (1.1)$$

where Y_t is R^d -valued, Z_t is $R^{d \times k}$ -valued, terminal value ξ is R^d -valued and $\mathcal{F}_T$ measurable. The generator function $f : [0, T] \times \Omega \times R^d \times R^{d \times k} \rightarrow R^d$ is of quadratic growth and W is a standard k -dimensional Brownian motion defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{P})$ where $(\mathcal{F}_t)$ is the Brownian filtration.

BSDE with quadratic growth can be used to solve problems such as utility maximization with exponential utility function. They were studied by Kobylanski [1] and then extended by others for example [2][3][4] and etc. More precisely in 2000, by using the monotonicity method adopted from PDE theory, Kobylanski [1] solved a class of one dimensional BSDEs with generator function being of quadratic growth in Z . This particular class of quadratic BSDEs with unbound terminal values were further studied by Briand and Hu [2][3] and Delbaen, Hu and Richou [4]. In 2013, Barrieu and El Karoui [5] adopted a different approach to prove the existence under conditions similar to those of Briand and Hu [2], while Briand and Elie [6] gave a concise study for the case when the terminal value ξ is bounded. The method used in the present paper to get the main result was partially inspired by the method used in Tevzadze [9] for solving existence of solution to a quadratic BSDE driven by a continuous martingale with bounded terminal values. The case of multidimensional quadratic BSDEs seems significantly more difficult than that of Lipschitz BSDEs, and the methods used in literature are often quite involved, and up until now the results about quadratic BSDEs are mostly only for the one-dimensional case, and they heavily rely on comparison theorems. Recently, based

^{*}Exeter College, Turl Street, Oxford OX1 3DP. Research supported partially by ERC grant ESig ID 291244. Email: qianz@maths.ox.ac.uk

[†]Mathematical Institute, University of Oxford, Oxford OX2 6GG. Email: yimin.yang@maths.ox.ac.uk

[‡]School of Finance and Statistics, East China Normal University, 500 Dongchuan Road, Shanghai 200241. Email: sjwu@stat.ecnu.edu.cn

on a result for one-dimensional BSDEs in Briand and Hu [2], Hu and Tang [12] proved the existence and uniqueness of solution to a multidimensional BSDE with diagonal quadratic generator assuming that each component f^i of the generator f depends only on the i -th row of the matrix variable Z in the BSDE (1.1).

In this paper we study a multidimensional quadratic BSDE with a particular class of product generators and give a result of existence of solution on a suitable Banach space under some constraints on parameters. The corresponding PDEs however have significance in fluid dynamics and in fact they are simplified version of fluid equations. The paper is organized as follows. In Section 2, we point out the particular type of multidimensional quadratic BSDEs that we study and list the assumptions we work under. In Section 3, we firstly use properties of BMO martingales, Girsanov's theorem and predictable representation property to show that the usual iteration method is also well defined for our problem. Then we show that a contraction map can be found on a suitable complete metric space on a fixed small time interval under some constraints on parameters, which gives a result of existence of solution to our BSDE on the whole time interval by pasting time together.

2 Definitions and assumptions

Let us begin with a few notations and definitions which are nevertheless standard in BSDE literature as follows:

- $\|y\| = \sqrt{y^T y}$ for $y \in \mathbb{R}^d$ and $\|z\| = \sqrt{\text{Tr}(zz^T)}$ for $z \in \mathbb{R}^{d \times k}$ denote the Euclidean norms.
- $E^{\mathcal{F}_t}[\cdot] := E[\cdot | \mathcal{F}_t]$
- W is a standard k -dimensional Brownian motion defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{P})$ and $(\mathcal{F}_t)_{t \in [0, T]}$ is the Brownian filtration.
- $\mathcal{M}^2(\mathbb{R}^d)$ and $\mathcal{M}^2(\mathbb{R}^{d \times k})$ denote respectively the Banach spaces of progressively measurable processes Y and Z such that $\|Y\|_{\mathcal{M}^2}^2 = E \int_0^T \|Y_t\|^2 dt < \infty$ and $\|Z\|_{\mathcal{M}^2}^2 = E \int_0^T \|Z_t\|^2 dt < \infty$.
- $\mathcal{S}^\infty(\mathbb{R}^d)$ denotes the Banach space of bounded progressively measurable processes Y .
- $\mathcal{S}_{C_1}^\infty(\mathbb{R}^d)$ denotes the Banach space of bounded progressively measurable processes Y such that $\|Y\|_{\mathcal{S}^\infty} \leq C_1$ and C_1 is a non negative constant.
- $\mathcal{B}(\mathbb{R}^{d \times k})$ denotes the space of progressively measurable processes Z such that $\int Z_s dW_s$ is a BMO martingale and define $\|Z\|_{\mathcal{B}} = \left\| \int Z_s dW_s \right\|_{BMO}$. Then $(\mathcal{B}, \|\cdot\|_{\mathcal{B}})$ is a Banach space due to the fact that the space of BMO martingales null at zero is Banach and the definition of stochastic integral.
- $\mathcal{B}_{\sqrt{R}}(\mathbb{R}^{d \times k}) = \{Z \in \mathcal{B}(\mathbb{R}^{d \times k}) : \|Z\|_{\mathcal{B}} \leq \sqrt{R}\}$ with R being a positive constant, which denotes a closed subspace of $\mathcal{B}(\mathbb{R}^{d \times k})$.

We consider the following BSDE:

$$\begin{cases} dY_t &= Z_t f(Y_t, Z_t) dt + Z_t dW_t \\ Y_T &= \xi \end{cases} \quad (2.1)$$

where Y_t is $\mathbb{R}^d$ -valued, Z_t is $\mathbb{R}^{d \times k}$ -valued, f is $\mathbb{R}^k$ -valued and ξ is $\mathbb{R}^d$ -valued and $\mathcal{F}_T$ measurable, which should be interpreted as a stochastic integral equation (1.1).

We make the following assumptions:

1. $\|\xi\| \leq C_1$ for some constant $C_1 > 0$, i.e. ξ is bounded.

2. f satisfies the Lipschitz condition and has a linear growth:

$$\begin{aligned} \|f(y_1, z_1) - f(y_2, z_2)\| &\leq C_2 \|y_1 - y_2\| + C_3 \|z_1 - z_2\|, \\ \|f(y_1, z_1)\| &\leq C_2 \|y_1\| + C_3 \|z_1\| + C_4, \end{aligned}$$

for any $y_1, y_2 \in \mathbb{R}^d$ and $z_1, z_2 \in \mathbb{R}^{d \times k}$, so that $zf(y, z)$ has quadratic growth in z . C_2, C_3, C_4 are non negative constants. And $f(Y, Z)$ is progressively measurable when (Y, Z) is progressively measurable.

By a solution to (2.1), we mean a pair of stochastic processes (Y, Z) on $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{P})$, where $Y = (Y_t) \in \mathcal{S}^\infty(\mathbb{R}^d)$ and $Z = (Z_t) \in \mathcal{B}(\mathbb{R}^{d \times k})$. Moreover $Z \in \mathcal{B}(\mathbb{R}^{d \times k})$ implies that $Z \in \mathcal{M}^2(\mathbb{R}^{d \times k})$ due to the fact that

$$\|Z\|_{\mathcal{M}^2}^2 = E \left(\int_0^T \|Z_s\|^2 ds \right) = E \left\| \int_0^T Z_s dW_s \right\|^2 \leq \left\| \int Z_s dW_s \right\|_{BMO}^2. \quad (2.2)$$

The following properties about BMO martingales are well known. If $\int Z_s dW_s$ is a BMO martingale, then

$$E^{\mathcal{F}_t} \left(\int_t^T \|Z\|^2 ds \right) \leq \left\| \int Z_s dW_s \right\|_{BMO}^2 = \|Z\|_{\mathcal{B}}^2, \forall t \in [0, T]$$

and if $E^{\mathcal{F}_t} \left(\int_t^T \|\tilde{Z}\|^2 ds \right) \leq N$, for every $t \in [0, T]$ where N is a non negative constant, then $\int \tilde{Z}_s dW_s$ is a BMO martingale and

$$\|\tilde{Z}\|_{\mathcal{B}}^2 = \left\| \int \tilde{Z}_s dW_s \right\|_{BMO}^2 \leq N,$$

see Kazamaki [10] for details.

3 Existence of solution

We will use the iteration method. Let $\mathcal{S}_{C_1}^\infty \times \mathcal{B}$ denote, for simplicity, the Banach space $\mathcal{S}_{C_1}^\infty(\mathbb{R}^d) \times \mathcal{B}(\mathbb{R}^{d \times k})$. Suppose that $(Y, Z) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}$ and f and ξ satisfy the above assumptions, then we have, by the linear growth of f , boundedness of Y and properties of the BMO martingale $\int Z_s dW_s$, that $\int f(Y_s, Z_s)^T dW_s$ is also a BMO martingale, which in turn implies that the stochastic exponential of $-\int f(Y_s, Z_s)^T dW_s$ is a martingale on $[0, T]$. Hence we define a probability measure $\mathbb{Q}$ by

$$\frac{d\mathbb{Q}}{d\mathbb{P}} \Big|_{\mathcal{F}_T} = \mathcal{E} \left(- \int f(Y_s, Z_s)^T dW_s \right)_T \quad (3.1)$$

and define

$$dW_t^{\mathbb{Q}} = dW_t + f(Y_t, Z_t) dt. \quad (3.2)$$

Then $W^{\mathbb{Q}}$ is a standard k -dimensional Brownian motion under probability measure $\mathbb{Q}$. The lemma below about continuous martingale representation is well known and its proof may be found in Cohen and Elliott [15].

Lemma 1. *Suppose M is a d -dimensional continuous local martingale under $\mathbb{Q}$ with $\mathbb{Q}$ defined as above, then there exists a unique predictable process H such that $M - M_0 = \int H_s dW_s^{\mathbb{Q}}$.*

Proof. Since M is a continuous semi-martingale under $\mathbb{P}$, $M - M_0 = N + A$ where N is a continuous local martingale null at 0 under $\mathbb{P}$ and A is a finite variation process. By the martingale representation theorem applying to N , we have that $N = \int H_s dW_s$ for some predictable process H . Thus

$M - M_0 - \int H_s dW_s^{\mathbb{Q}} = -\int H_s f(Y_s, Z_s) ds + A$ by equation (3.2), which implies that the continuous $\mathbb{Q}$ -local martingale $M - M_0 - \int H_s dW_s^{\mathbb{Q}}$ is of finite variation and null at 0. Thus we have that $M - M_0 = \int H_s dW_s^{\mathbb{Q}}$. Uniqueness can be proved in the usual way. $\square$

Let $\delta \in (0, 1)$, and consider time interval $[T - \delta T, T]$. Let $(Y, Z) \in \mathcal{S}_{C_1}^{\infty} \times \mathcal{B}$ but with duration $[T - \delta T, T]$.

Since $\|\xi\| \leq C_1$, $\tilde{Y}_t = E_{\mathbb{Q}}^{\mathcal{F}_t}[\xi]$ is a continuous martingale under $\mathbb{Q}$ on $[T - \delta T, T]$. Thus by Lemma 1, there exists a unique predictable process $\tilde{Z}$ on $[T - \delta T, T]$ such that

$$\begin{cases} d\tilde{Y}_t &= \tilde{Z}_t f(Y_t, Z_t) dt + \tilde{Z}_t dW_t \\ \tilde{Y}_T &= \xi \end{cases} \quad (3.3)$$

with $\|\tilde{Y}\|_{\mathcal{S}^{\infty}} \leq C_1$, which means that $\tilde{Y} \in \mathcal{S}_{C_1}^{\infty}(R^d)$. We prove the following proposition.

Proposition 2. *If $C_1 C_3 < e^{-\frac{1}{2}}$ where $e^{-\frac{1}{2}}$ is just a universal constant, and it does not imply that it is optimal. Then there is a non negative constant C_6 depending on C_1, C_2, C_3, C_4 and δT such that*

$$\|\tilde{Z}\|_{\mathcal{B}}^2 \leq C_6 + \frac{1}{2} \|Z\|_{\mathcal{B}}^2$$

for any pairs (Y, Z) and $(\tilde{Y}, \tilde{Z})$ on $[T - \delta T, T]$ defined by BSDE (3.3).

Proof. Consider $\varphi(t) = e^{Kt}$ where K is a positive constant to be determined later. Let $\eta_t = \varphi(\|\tilde{Y}_t\|^2)$. Then by Ito's formula we have

$$\begin{aligned} d\|\tilde{Y}\|^2 &= 2\sum_i \tilde{Y}^i d\tilde{Y}^i + \|\tilde{Z}\|^2 dt \\ &= 2\sum_{i,j} f_j(Y, Z) \tilde{Y}^i \tilde{Z}^{i,j} dt + \|\tilde{Z}\|^2 dt + 2\sum_{i,j} \tilde{Y}^i \tilde{Z}^{i,j} dW^j, \end{aligned}$$

and

$$\begin{aligned} d\eta &= K\eta d\|\tilde{Y}\|^2 + 2K^2\eta \sum_j \left| \sum_i \tilde{Y}^i \tilde{Z}^{i,j} \right|^2 dt \\ &= K\eta \left[2\sum_{i,j} f_j(Y, Z) \tilde{Y}^i \tilde{Z}^{i,j} dt + \|\tilde{Z}\|^2 dt \right] + 2K^2\eta \sum_j \left| \sum_i \tilde{Y}^i \tilde{Z}^{i,j} \right|^2 dt \\ &\quad + 2K\eta \sum_{i,j} \tilde{Y}^i \tilde{Z}^{i,j} dW^j. \end{aligned}$$

Set vector $\tilde{U}$ with $\tilde{U}^j = \sum_i \tilde{Y}^i \tilde{Z}^{i,j}$, so the previous equation can be written as

$$d\eta = K\eta \left[2\tilde{U}^T f(Y, Z) + \|\tilde{Z}\|^2 + 2K\|\tilde{U}\|^2 \right] dt + 2K\eta \tilde{U}^T dW.$$

Integrating the equality above from t to T , we obtain

$$\eta_t = \eta_T - K \int_t^T \eta \left[2\tilde{U}^T f(Y, Z) + \|\tilde{Z}\|^2 + 2K\|\tilde{U}\|^2 \right] ds - 2K \int_t^T \eta \tilde{U}^T dW.$$

Take the conditional expectation with respect to $\mathcal{F}_t$ to get

$$\eta_t = E^{\mathcal{F}_t}(\eta_T) - KE^{\mathcal{F}_t} \left(\int_t^T \eta \left[2\tilde{U}^T f(Y, Z) + \|\tilde{Z}\|^2 + 2K\|\tilde{U}\|^2 \right] ds \right).$$

Next applying Cauchy-Schwartz inequality, the linear growth condition of f and the bound of Y , we deduce that

$$\begin{aligned}\eta_t &\leq E^{\mathcal{F}_t}(\eta_T) - KE^{\mathcal{F}_t} \left(\int_t^T \eta \left[-2(C_1C_2 + C_3\|Z\| + C_4)\|\tilde{U}\| + \|\tilde{Z}\|^2 + 2K\|\tilde{U}\|^2 \right] ds \right) \\ &= E^{\mathcal{F}_t}(\eta_T) + 2(C_1C_2 + C_4)KE^{\mathcal{F}_t} \left(\int_t^T \eta \|\tilde{U}\| ds \right) \\ &\quad + 2KC_3E^{\mathcal{F}_t} \left(\int_t^T \eta \|Z\| \|\tilde{U}\| ds \right) - KE^{\mathcal{F}_t} \left(\int_t^T \eta \left[\|\tilde{Z}\|^2 + 2K\|\tilde{U}\|^2 \right] ds \right).\end{aligned}$$

Then by applying the inequalities with $\alpha, \beta > 0$

$$2\|\tilde{U}\| \leq \alpha + \frac{1}{\alpha}\|\tilde{U}\|^2$$

and

$$2\|Z\|\|\tilde{U}\| \leq \beta\|Z\|^2 + \frac{1}{\beta}\|\tilde{U}\|^2,$$

we get that

$$\begin{aligned}\eta_t &\leq E^{\mathcal{F}_t}(\eta_T) + (C_1C_2 + C_4)KE^{\mathcal{F}_t} \left(\int_t^T \alpha \eta ds \right) \\ &\quad - KE^{\mathcal{F}_t} \left(\int_t^T \eta \|\tilde{Z}\|^2 ds \right) + KC_3E^{\mathcal{F}_t} \left(\int_t^T \eta \beta \|Z\|^2 ds \right) \\ &\quad - KE^{\mathcal{F}_t} \left(\int_t^T \eta \left[2K - \frac{C_3}{\beta} - \frac{(C_1C_2 + C_4)}{\alpha} \right] \|\tilde{U}\|^2 ds \right).\end{aligned}$$

It follows that

$$\begin{aligned}E^{\mathcal{F}_t} \left(\int_t^T \eta \|\tilde{Z}\|^2 ds \right) &\leq \frac{1}{K}E^{\mathcal{F}_t}(\eta_T - \eta_t) + (C_1C_2 + C_4)E^{\mathcal{F}_t} \left(\int_t^T \alpha \eta ds \right) \\ &\quad + C_3\beta E^{\mathcal{F}_t} \left(\int_t^T \eta \|Z\|^2 ds \right) \\ &\quad - E^{\mathcal{F}_t} \left(\int_t^T \eta \left[2K - \frac{C_3}{\beta} - \frac{(C_1C_2 + C_4)}{\alpha} \right] \|\tilde{U}\|^2 ds \right).\end{aligned}\quad (3.4)$$

Since $\|\tilde{Y}\|_{\mathcal{G}^\infty} \leq C_1$ we deduce that $1 \leq \eta \leq e^{KC_1^2}$. We may choose constants such that

$$2K - \frac{C_3}{\beta} - \frac{(C_1C_2 + C_4)}{\alpha} \geq 0 \quad (3.5)$$

and

$$C_3\beta e^{KC_1^2} = \frac{1}{2}. \quad (3.6)$$

In order to do this, it requires that

$$K - C_3^2 e^{KC_1^2} > 0$$

which means that

$$Ke^{-1} - C_3^2 e^{KC_1^2 - 1} > 0$$

which is possible only if $C_1C_3 < e^{-\frac{1}{2}}$. When $C_1C_3 < e^{-\frac{1}{2}}$ by considering

$$KC_1^2 - C_1^2C_3^2 e^{KC_1^2} > 0$$

we can choose

$$K = -\frac{2}{C_1^2} \ln(C_1 C_3)$$

which implies that

$$K - C_3^2 e^{KC_1^2} > 0.$$

Then we can deduce from the previous inequality (3.4) that

$$\begin{aligned} E^{\mathcal{F}_t} \left(\int_t^T \|\tilde{Z}\|^2 ds \right) &\leq \frac{1}{K} e^{KC_1^2} + \alpha e^{KC_1^2} (C_1 C_2 + C_4) \delta T \\ &\quad + C_3 \beta e^{KC_1^2} E^{\mathcal{F}_t} \left(\int_t^T \|Z\|^2 ds \right) \end{aligned}$$

for all $t \in [T - \delta T, T]$. Using the properties of BMO martingales we may deduce that

$$\|\tilde{Z}\|_{\mathcal{B}}^2 \leq C_6 + \frac{1}{2} \|Z\|_{\mathcal{B}}^2$$

where

$$\begin{aligned} C_6 &= e^{KC_1^2} \left[\frac{1}{K} + \alpha (C_1 C_2 + C_4) \delta T \right] \\ &= \frac{1}{C_1^2 C_3^2} \left[\frac{C_1^2}{-2 \ln(C_1 C_3)} + \alpha (C_1 C_2 + C_4) \delta T \right]. \end{aligned} \quad (3.7)$$

□

The above proposition implies that we get a pair $(\tilde{Y}, \tilde{Z}) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}$ on $[T - \delta T, T]$, when $C_1 C_3 < e^{-\frac{1}{2}}$. In this case we define the pair $(\tilde{Y}, \tilde{Z}) = \Phi(Y, Z)$ on $[T - \delta T, T]$ and $\Phi : \mathcal{S}_{C_1}^\infty \times \mathcal{B} \rightarrow \mathcal{S}_{C_1}^\infty \times \mathcal{B}$ is well defined.

In order to get a result about the global existence of solution, we firstly consider the time interval $[T - \delta T, T]$ for some $\delta \in (0, 1)$ and try to find a contraction map on a closed subspace of $\mathcal{S}_{C_1}^\infty \times \mathcal{B}$. This approach is inspired by the method used in Tevzadze [9]. Then by working backwards with respect to time intervals of length δT , we can get our result by pasting time together.

Theorem 3. *Under the above assumptions on f and ξ . If $C_1 C_3 < e^{-324}$, then for any terminal time T , there exists a positive constant $\tilde{R} = \lceil \frac{1}{\delta} \rceil R$ with $R = 2C_6$ and some fixed constant $\delta \in (0, 1)$ such that the BSDE*

$$\begin{cases} dY_t &= Z_t f(Y_t, Z_t) dt + Z_t dW_t \\ Y_T &= \xi \end{cases} \quad (3.8)$$

has a pair of solution $(Y, Z) \in \mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{\tilde{R}}}(R^{d \times k})$ on $[0, T]$.

Proof. Let $\lambda \in (0, 1)$ be a positive constant to be determined later. We firstly consider the time interval $[T - \delta T, T]$ as above and assume that

$$C_1 C_3 < e^{-\frac{4}{\lambda^2}} \quad (3.9)$$

which implies that $C_1 C_3 < e^{-\frac{1}{2}}$ and we set constant R to be

$$R = 2C_6. \quad (3.10)$$

We may have the following by choosing δ small enough.

$$2(C_1C_2 + C_4)\sqrt{\delta T} \vee 2C_2\sqrt{\delta T}\sqrt{R} \vee 2C_3\sqrt{R} \leq \lambda. \quad (3.11)$$

We can do this because we have condition (3.9) and in equation (3.7):

$$C_6 = \frac{1}{C_1^2 C_3^2} \left[\frac{C_1^2}{-2\ln(C_1 C_3)} + \alpha(C_1 C_2 + C_4) \delta T \right]$$

which implies that

$$C_3^2 C_6 = \frac{1}{C_1^2} \left[\frac{C_1^2}{-2\ln(C_1 C_3)} + \alpha(C_1 C_2 + C_4) \delta T \right],$$

where α is determined by inequality (3.5):

$$2K - \frac{C_3}{\beta} - \frac{(C_1 C_2 + C_4)}{\alpha} \geq 0$$

which can be achieved when $C_1 C_3 < e^{-\frac{1}{2}}$.

Let $\mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ denote the Banach space $\mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{R}}(R^{d \times k})$. Since $C_1 C_3 < e^{-\frac{1}{2}}$ which is due to condition (3.9), we have $\Phi : \mathcal{S}_{C_1}^\infty \times \mathcal{B} \rightarrow \mathcal{S}_{C_1}^\infty \times \mathcal{B}$ as defined above. Then for any pair $(Y, Z) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ we can get $(\tilde{Y}, \tilde{Z}) = \Phi(Y, Z)$ with $(\tilde{Y}, \tilde{Z}) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}$. By Proposition 2 we have that

$$\|\tilde{Z}\|_{\mathcal{B}}^2 \leq C_6 + \frac{1}{2} \|Z\|_{\mathcal{B}}^2.$$

Together with condition (3.10) we get that

$$\|\tilde{Z}\|_{\mathcal{B}}^2 \leq \frac{R}{2} + \frac{1}{2} \|Z\|_{\mathcal{B}}^2 \leq R,$$

which implies that $(\tilde{Y}, \tilde{Z}) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$. So that $\Phi : \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}} \rightarrow \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ is well defined.

For any $(Y^1, Z^1), (Y^2, Z^2) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$, we set $(\tilde{Y}^1, \tilde{Z}^1) = \Phi(Y^1, Z^1)$ and $(\tilde{Y}^2, \tilde{Z}^2) = \Phi(Y^2, Z^2)$. So we have $(\tilde{Y}^1, \tilde{Z}^1), (\tilde{Y}^2, \tilde{Z}^2) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$. Then by setting

$$\Delta = Y^1 - Y^2, \tilde{\Delta} = \tilde{Y}^1 - \tilde{Y}^2, \Lambda = Z^1 - Z^2, \tilde{\Lambda} = \tilde{Z}^1 - \tilde{Z}^2,$$

we get that $(\Delta, \Lambda), (\tilde{\Delta}, \tilde{\Lambda}) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ with $\tilde{\Delta}_T = 0$ and we also get

$$\begin{aligned} d\tilde{\Delta}^i &= \sum_j \tilde{\Lambda}^{i,j} dW^j + \sum_j f_j(Y^1, Z^1) \tilde{\Lambda}^{i,j} dt \\ &\quad + \sum_j [f_j(Y^1, Z^1) - f_j(Y^2, Z^2)] \tilde{Z}^{i,j} dt. \end{aligned}$$

Then by Ito's formula we have

$$\begin{aligned} d\|\tilde{\Delta}\|^2 &= 2\vartheta^T f(Y^1, Z^1) dt + 2\rho^T [f(Y^1, Z^1) - f(Y^2, Z^2)] dt \\ &\quad + \|\tilde{\Lambda}\|^2 dt + 2\vartheta^T dW, \end{aligned} \quad (3.12)$$

where the components of vectors ϑ and ρ are defined as

$$\vartheta^j = \sum_i \tilde{\Delta}^i \tilde{\Lambda}^{i,j}, \quad \rho^j = \sum_i \tilde{\Delta}^i \tilde{Z}^{i,j},$$

and $\int \vartheta^T dW$ is a martingale. Then by taking conditional expectation we get

$$\begin{aligned} \|\tilde{\Delta}_t\|^2 + E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Lambda}\|^2 ds \right] &= -2E^{\mathcal{F}_t} \left[\int_t^T \vartheta^T f(Y^1, Z^1) ds \right] \\ &\quad - 2E^{\mathcal{F}_t} \left[\int_t^T \rho^T [f(Y^1, Z^1) - f(Y^2, Z^2)] ds \right], \end{aligned}$$

which implies that

$$\begin{aligned} \|\tilde{\Delta}_t\|^2 + E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Lambda}\|^2 ds \right] &\leq 2E^{\mathcal{F}_t} \left[\int_t^T \|\vartheta\| \|f(Y^1, Z^1)\| ds \right] \\ &\quad + 2E^{\mathcal{F}_t} \left[\int_t^T \|\rho\| \|f(Y^1, Z^1) - f(Y^2, Z^2)\| ds \right]. \end{aligned}$$

Together with the definition of ϑ and ρ , we obtain from the inequality above that

$$\begin{aligned} \|\tilde{\Delta}_t\|^2 + E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Lambda}\|^2 ds \right] &\leq 2E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Delta}\| \|\tilde{\Lambda}\| \|f(Y^1, Z^1)\| ds \right] \\ &\quad + 2E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Delta}\| \|\tilde{Z}^2\| \|f(Y^1, Z^1) - f(Y^2, Z^2)\| ds \right] \end{aligned}$$

for all $t \in [T - \delta T, T]$.

Now by using the assumptions on f , we conclude that

$$\begin{aligned} \|\tilde{\Delta}_t\|^2 + E^{\mathcal{F}_t} \left[\int_t^T \|\tilde{\Lambda}_s\|^2 ds \right] &\leq 2E^{\mathcal{F}_t} \left[\int_t^T [(C_1 C_2 + C_4) + C_3 \|Z_s^1\|] \|\tilde{\Delta}_s\| \|\tilde{\Lambda}_s\| ds \right] \\ &\quad + 2E^{\mathcal{F}_t} \left[\int_t^T (C_2 \|\Delta_s\| + C_3 \|\Lambda_s\|) \|\tilde{\Delta}_s\| \|\tilde{Z}_s^2\| ds \right] \\ &\leq 2(C_1 C_2 + C_4) \sqrt{\delta T} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} + 2C_3 \|Z^1\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} \\ &\quad + 2C_2 \sqrt{\delta T} \|\tilde{Z}^2\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Delta\|_{\mathcal{J}^\infty} + 2C_3 \|\tilde{Z}^2\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Lambda\|_{\mathcal{B}}, \end{aligned}$$

from which we deduce that

$$\begin{aligned} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} &\leq 2(C_1 C_2 + C_4) \sqrt{\delta T} \|\tilde{\Lambda}\|_{\mathcal{B}} + 2C_3 \|Z^1\|_{\mathcal{B}} \|\tilde{\Lambda}\|_{\mathcal{B}} \\ &\quad + 2C_2 \sqrt{\delta T} \|\tilde{Z}^2\|_{\mathcal{B}} \|\Delta\|_{\mathcal{J}^\infty} + 2C_3 \|\tilde{Z}^2\|_{\mathcal{B}} \|\Lambda\|_{\mathcal{B}}, \end{aligned} \tag{3.13}$$

and

$$\begin{aligned} \|\tilde{\Lambda}\|_{\mathcal{B}}^2 &\leq 2(C_1 C_2 + C_4) \sqrt{\delta T} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} + 2C_3 \|Z^1\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} \\ &\quad + 2C_2 \sqrt{\delta T} \|\tilde{Z}^2\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Delta\|_{\mathcal{J}^\infty} + 2C_3 \|\tilde{Z}^2\|_{\mathcal{B}} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Lambda\|_{\mathcal{B}}. \end{aligned} \tag{3.14}$$

Thus we have that

$$\begin{aligned} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} &\leq 2(C_1 C_2 + C_4) \sqrt{\delta T} \|\tilde{\Lambda}\|_{\mathcal{B}} + 2C_3 \sqrt{R} \|\tilde{\Lambda}\|_{\mathcal{B}} \\ &\quad + 2C_2 \sqrt{\delta T} \sqrt{R} \|\Delta\|_{\mathcal{J}^\infty} + 2C_3 \sqrt{R} \|\Lambda\|_{\mathcal{B}}, \end{aligned} \tag{3.15}$$

and

$$\begin{aligned} \|\tilde{\Lambda}\|_{\mathcal{B}}^2 &\leq 2(C_1 C_2 + C_4) \sqrt{\delta T} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} + 2C_3 \sqrt{R} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\tilde{\Lambda}\|_{\mathcal{B}} \\ &\quad + 2C_2 \sqrt{\delta T} \sqrt{R} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Delta\|_{\mathcal{J}^\infty} + 2C_3 \sqrt{R} \|\tilde{\Delta}\|_{\mathcal{J}^\infty} \|\Lambda\|_{\mathcal{B}}. \end{aligned} \tag{3.16}$$

Then by condition (3.11) we get that

$$\|\tilde{\Delta}\|_{\mathcal{S}^\infty} \leq \lambda (2\|\tilde{\Lambda}\|_{\mathcal{B}} + \|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}) \quad (3.17)$$

and

$$\|\tilde{\Lambda}\|_{\mathcal{B}}^2 \leq \lambda \|\tilde{\Delta}\|_{\mathcal{S}^\infty} (2\|\tilde{\Lambda}\|_{\mathcal{B}} + \|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}). \quad (3.18)$$

So by substituting $\|\tilde{\Delta}\|_{\mathcal{S}^\infty}$ in (3.18) with (3.17) we have

$$\|\tilde{\Lambda}\|_{\mathcal{B}}^2 \leq 8\lambda^2 (\|\tilde{\Lambda}\|_{\mathcal{B}}^2 + \|\Delta\|_{\mathcal{S}^\infty}^2 + \|\Lambda\|_{\mathcal{B}}^2),$$

which implies that

$$\|\tilde{\Lambda}\|_{\mathcal{B}} \leq 3\lambda (\|\tilde{\Lambda}\|_{\mathcal{B}} + \|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}).$$

By combining with (3.17) we deduce that

$$\|\tilde{\Delta}\|_{\mathcal{S}^\infty} + \|\tilde{\Lambda}\|_{\mathcal{B}} \leq 5\lambda \|\tilde{\Lambda}\|_{\mathcal{B}} + 4\lambda (\|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}).$$

If $\lambda < \frac{1}{5}$ we have that

$$\|\tilde{\Delta}\|_{\mathcal{S}^\infty} + \|\tilde{\Lambda}\|_{\mathcal{B}} \leq \frac{4\lambda}{(1-5\lambda)} (\|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}). \quad (3.19)$$

If we can choose $\lambda < \frac{1}{9}$ so that $\frac{4\lambda}{(1-5\lambda)} < 1$. Then $\Phi : \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}} \rightarrow \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ is a contraction map. By the Banach's fixed point theorem, we deduce that there exists a unique pair of solution $(Y, Z) \in \mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ on the time interval $[T - \delta T, T]$ to the BSDE (3.8) restricted on $\mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$. Therefore what left to be shown is that there exists $\lambda < \frac{1}{9}$ such that assumption (3.9) holds, and this

can be achieved when $C_1 C_3 < e^{-\frac{4}{(\frac{1}{9})^2}} = e^{-324}$.

We then consider the time interval $[T - 2\delta T, T - \delta T]$ if $T - 2\delta T > 0$ and $[0, T - \delta T]$ otherwise and set terminal value ξ at time $T - \delta T$ to be $Y_{T-\delta T}$ which is the initial value of the solution Y solved above on the time interval $[T - \delta T, T]$. Then by using the above same method, we get a unique pair of solution in $\mathcal{S}_{C_1}^\infty \times \mathcal{B}_{\sqrt{R}}$ on the time interval $[T - 2\delta T, T - \delta T]$ to the BSDE (3.8). By repeating this procedure backwards and pasting the solutions on all the time intervals together we get a pair of solution $(Y, Z) \in \mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{R}}(R^{d \times k})$ with $\tilde{R} = \lceil \frac{1}{\delta} \rceil R$ on the time interval $[0, T]$ to the BSDE (3.8).

If δ and λ satisfy the condition that $\sqrt{\lceil \frac{1}{\delta} \rceil} \lambda < \frac{1}{9}$, which may be achievable when T and $C_1 C_3$ are small enough. Then the pair of solution $(Y, Z) \in \mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{R}}(R^{d \times k})$ on the time interval $[0, T]$ to the BSDE (3.8) is unique. This uniqueness of the solution can be proved as follows. Suppose there exist two pairs of solutions $(Y^1, Z^1), (Y^2, Z^2) \in \mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{R}}(R^{d \times k})$ on the time interval $[0, T]$ to the BSDE (3.8). By setting

$$\Delta = Y^1 - Y^2, \Lambda = Z^1 - Z^2,$$

we get that $(\Delta, \Lambda) \in \mathcal{S}_{C_1}^\infty(R^d) \times \mathcal{B}_{\sqrt{R}}(R^{d \times k})$ with $\Delta_T = 0$. Then on the time interval $[T - \delta T, T]$ by repeating the procedure starting from equation (3.12), we obtain an inequality which is similar to inequality (3.19) as follows:

$$\|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}} \leq \frac{4\sqrt{\lceil \frac{1}{\delta} \rceil} \lambda}{(1 - 5\sqrt{\lceil \frac{1}{\delta} \rceil} \lambda)} (\|\Delta\|_{\mathcal{S}^\infty} + \|\Lambda\|_{\mathcal{B}}).$$

Since $\lambda < \frac{1}{9\sqrt{\lceil \frac{1}{8} \rceil}}$ so that $\frac{4\sqrt{\lceil \frac{1}{8} \rceil}\lambda}{(1-5\sqrt{\lceil \frac{1}{8} \rceil}\lambda)} < 1$, we deduce that $\|\Delta\|_{\mathcal{S}^\infty} = 0$ and $\|\Lambda\|_{\mathcal{B}} = 0$. Thus (Y^1, Z^1) equals (Y^2, Z^2) on the time interval $[T - \delta T, T]$, in particular $Y_{T-\delta T}^1$ equals $Y_{T-\delta T}^2$. We then consider the time interval $[T - 2\delta T, T - \delta T]$ if $T - 2\delta T > 0$ and $[0, T - \delta T]$ otherwise, and terminal values at time $T - \delta T$ are $Y_{T-\delta T}^1$ and $Y_{T-\delta T}^2$ respectively for the two solutions. By using the same procedure starting from equation (3.12), we deduce that (Y^1, Z^1) equals (Y^2, Z^2) on the time interval $[T - 2\delta T, T - \delta T]$ as well. Thus by repeating this procedure backwards, we conclude that (Y^1, Z^1) equals (Y^2, Z^2) on the time interval $[0, T]$. $\square$

Remark 4. Theorem 3 says that, given parameters C_2, C_3, C_4 , if the bound C_1 of the terminal value is small enough then there exists a pair of solution $(Y, Z) \in \mathcal{S}^\infty(R^d) \times \mathcal{B}(R^{d \times k})$ on the time interval $[0, T]$ to the BSDE (3.8).

References

- [1] M.Kobylanski. Backward stochastic differential equations and partial differential equations with quadratic growth. The Annals of Probability, Vol.28, No. 2, 558–602 (2000).
- [2] P.Briand and Y.Hu. BSDE with quadratic growth and unbounded terminal value. Probability Theory and Related Fields, 136, No.4, 604–618 (2006).
- [3] P.Briand and Y.Hu. Quadratic BSDEs with convex generators and unbounded terminal conditions. Probability Theory and Related Fields, 141, 543–567 (2008).
- [4] F.Delbaen, Y.Hu and A.Richou. On the uniqueness of solutions to quadratic BSDEs with convex generators and unbounded terminal conditions. Probabilités et Statistiques, Vol.47, No.2, 559–574 (2011).
- [5] P.Barrieu and N.El Karoui. Monotone stability of quadratic semimartingales with applications to general quadratic BSDEs. The Annals of Probability, Vol 41, 1831–1863 (2013).
- [6] P.Briand and R.Elle. A simple constructive approach to quadratic BSDEs with or without delay. Stochastic Processes and their Applications, 123(8), 2921–2939 (2013).
- [7] J.P. Lepeltier and J. San Martin. Backward stochastic differential equations with continuous coefficients. Statistics & Probability Letters, 32, 425–430 (1997).
- [8] Mark H.A. Davis. Advances in Control, Communication Networks, and Transportation Systems: In Honor of Pravin Varaiya, chapter Martingale representation and all that. Systems and Control: Foundations and Applications. Birkhauser, (2005).
- [9] R.Tevzadze. Solvability of backward stochastic differential equations with quadratic growth. Stochastic Processes and their Applications, 118(3), 503–515 (2008).
- [10] N.Kazamaki. Continuous exponential martingales and BMO, Lecture Notes in Mathematics, 1579. Springer-Verlag, Berlin (1994).
- [11] G.Liang, A.Lionnet and Z.Qian. (2010). On Girsanov’s transform for backward stochastic differential equations. Preprint arXiv:1011.3228 (2010).
- [12] Y.Hu and S.Tang. Multi-Dimensional Backward Stochastic Differential Equations of Diagonally Quadratic Generators. Stochastic Processes and their Applications, 126(4), 1066–1086 (2016).

- [13] G.Liang, T.Lyons and Z.Qian. Backward stochastic dynamics on a filtered probability space. *Annals of probability*, 39(4), 1422-1448, (2011).
- [14] E.Pardoux and S.G.Peng. Adapted solution of a backward stochastic differential equation. *Systems & Control letters*, 14, 55-61 (1990).
- [15] S. N. Cohen and R. J. Elliott. *Stochastic Calculus and Applications*. Springer-Verlag (2015).